



Genome Sequences of *Frankineae* sp. Strain MT45 and *Jatrophihabitans* sp. Strain GAS493, Two Actinobacteria Isolated from Forest Soil

 Kristen M. DeAngelis,^a  Grace Pold^{b,c}

^aDepartment of Microbiology, University of Massachusetts, Amherst, Massachusetts, USA

^bSchool of Natural Science, Hampshire College, Amherst, Massachusetts, USA

^cDepartment of Natural Resources Management and Environmental Sciences, California Polytechnic State University, San Luis Obispo, California, USA

ABSTRACT *Frankiaceae* are bacterial endosymbionts that are also found free-living in soil. Here, we present the genome sequences of two novel bacterial members of the order *Frankiales*, class *Actinobacteria*, isolated from temperate terrestrial forest soils. The genomes for MT45 and GAS493 indicate a genetic capacity for carbohydrate degradation but not nitrogen fixation.

Frankiaceae are best known as spore-forming, nitrogen-fixing actinobacteria (1) capable of associating with actinorhizal plants (2), although these plants are not required for their growth (3). Because their role in free-living nitrogen fixation is still unclear, we isolated and sequenced two new taxa that are unclassified in the family *Frankiaceae* and are genomically distinct from known type strains (4).

Frankineae sp. strain MT45 and *Jatrophihabitans* sp. strain GAS493 were both isolated from soils collected from an even-aged mixed-hardwood forest stand at the Harvard Forest in central Massachusetts. MT45 was isolated from the organic-rich forest floor, while GAS493 was isolated from the mineral soil below. Isolates were cultivated aerobically on VL55 medium solidified with gellan gum (5) and amended with xylan for MT45 and mixed plant polymers xylan, xanthan, pectin, and carboxymethyl cellulose for GAS493.

Genomic DNA was extracted using the Qiagen Genomic-tip 500/G kit. SMRTbell libraries were constructed and sequenced on the PacBio RS platform at the U.S. Department of Energy (DOE) Joint Genome Institute (JGI) (6), generating 383,007 reads filtered to 122,552 subreads totaling 611.7 Mbp for MT45 and 506,493 reads filtered to 259,823 subreads totaling 914.4 Mbp for GAS493. The libraries had an N_{50} value of 4.229 Mb for MT45 and 4.88 Mb for GAS493. The raw reads were assembled using HGAP (v.2.3.0 p5; protocol v.2.3.0, RS HGAP Assembly.3 method, smrtpipe.py v1.87.139483) (7). The final assembly was one 4.229-Mbp scaffold with 70.0 \times input read coverage for MT45 and one 4.880-Mbp scaffold with 156.0 \times coverage for GAS493. CheckM (8) indicates that the MT45 and GAS493 genomes are 95.76% and 95.91% complete with 1.4% and 1.5% contamination, respectively. The MT45 and GAS493 genomes have GC contents of 67.3% and 66.94%, respectively.

Genes were identified using Prodigal (9), manually curated using GenePRIMP (10), and then translated to search the NCBI nonredundant database and the UniProt, TIGRFam, Pfam, KEGG, Clusters of Orthologous Groups (COG), and InterPro databases. rRNA genes were found by searches against models of the rRNA genes built from SILVA (11). Gene searches were completed within the JGI Integrated Microbial Genomes (IMG) platform (12) and KBase (13). Default parameters were used for all software unless otherwise noted.

Genomes were compared by average nucleotide identity (ANI) and using Mash

Citation DeAngelis KM, Pold G. 2020. Genome sequences of *Frankineae* sp. strain MT45 and *Jatrophihabitans* sp. strain GAS493, two actinobacteria isolated from forest soil. *Microbiol Resour Announc* 9:e00614-20. <https://doi.org/10.1128/MRA.00614-20>.

Editor Irene L. G. Newton, Indiana University, Bloomington

Copyright © 2020 DeAngelis and Pold. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Kristen M. DeAngelis, deangelis@microbio.umass.edu.

Received 27 May 2020

Accepted 28 August 2020

Published 17 September 2020

(14) in KBase (13), which uses a whole-genome nucleotide clustering method to define genome similarity. GAS493 and MT45 share 84% ANI and a Minhash D of 0.147, or $1 - \text{ANI}$ (14). The next closest relative to GAS493 and MT45 was a *Geodermatophilus* species, which had D values of 0.241 and 0.244, respectively. MT45 has two rRNA operons; the two 16S rRNA genes are identical in sequence and bear 99% identity to those of *Jatrophihabitans* sp. GAS493 and less than 97% to any other cultivated bacteria.

Carbohydrate utilization is observed among only some *Frankiaceae* (1), and both MT45 and GAS493 have dozens of genes annotated for carbohydrate activity (15). Both lack the canonical genes for nitrogen fixation, but they do have a NifU-like gene encoding an Fe-S cluster assembly protein with diverse functions in cell redox (16). Both genomes contain genes required for nitrogen assimilation through ammonium reduction, including an ammonia permease gene, which are observed for free-living diazotrophs (1). Although some *Frankia* spp. can fix nitrogen based on the acetylene reduction assay without NifH homologs detected by PCR amplification (17), the mechanism is unknown. Therefore, we propose a potential role for GAS493 and MT45 as nondiazotrophic polysaccharide degraders in soil.

Together, these genomes suggest multiple mechanisms for growth in soils where nutrients can be poor and sparsely available.

Data availability. Raw reads are available for download in the Sequence Read Archive under the accession no. [SRX2158410](https://www.ncbi.nlm.nih.gov/sra/SRX2158410) for *Frankineae* sp. MT45 and accession no. [SRX3048269](https://www.ncbi.nlm.nih.gov/sra/SRX3048269) for *Jatrophihabitans* sp. GAS493. Genome assemblies have been deposited in GenBank under the accession no. [LT629697](https://www.ncbi.nlm.nih.gov/genbank/LT629697) for *Frankineae* sp. MT45 and accession no. [LT907982](https://www.ncbi.nlm.nih.gov/genbank/LT907982) for *Jatrophihabitans* sp. GAS493.

ACKNOWLEDGMENTS

Funding for this project came from the U.S. Department of Energy (DOE) Joint Genome Institute, a DOE Office of Science User Facility, which is supported by the Office of Science of the DOE under contract DE-AC02-05CH11231. This work was also supported by the National Science Foundation under contract DEB-1749206 and the DOE Genomic Sciences Program under contract DE-SC0016590.

We thank Andrew Billings and Samantha Murphy for assistance in the isolation and extraction of genomic DNA (gDNA) of bacterial isolates.

REFERENCES

- Normand P, Orso S, Cournoyer B, Jeannin P, Chapelon C, Dawson J, Evtushenko L, Misra AK. 1996. Molecular phylogeny of the genus *Frankia* and related genera and emendation of the family Frankiaceae. *Int J Syst Evol Microbiol* 46:1–9. <https://doi.org/10.1099/00207713-46-1-1>.
- Benson DR, Silvester WB. 1993. Biology of *Frankia* strains, actinomycece symbionts of actinorhizal plants. *Microbiol Mol Biol Rev* 57:293–319. <https://doi.org/10.1128/MMBR.57.2.293-319.1993>.
- Smolander A, Sundman V. 1987. *Frankia* in acid soils of forests devoid of actinorhizal plants. *Physiol Plant* 70:297–303. <https://doi.org/10.1111/j.1399-3054.1987.tb06147.x>.
- Nouioui I, Ghodhbane-Gtari F, del Carmen Montero-Calasanz M, Göker M, Meier-Kolthoff JP, Schumann P, Rohde M, Goodfellow M, Fernandez MP, Normand P, Tisa LS, Klenk H-P, Gtari M. 2016. Proposal of a type strain for *Frankia alni* (Woronin 1866) Von Tubeuf 1895, emended description of *Frankia alni*, and recognition of *Frankia casuarinae* sp. nov. and *Frankia elaeagni* sp. nov. *Int J Syst Evol Microbiol* 66:5201–5210. <https://doi.org/10.1099/ijsem.0.001496>.
- Eichorst SA, Kuske CR, Schmidt TM. 2011. Influence of plant polymers on the distribution and cultivation of bacteria in the phylum Acidobacteria. *Appl Environ Microbiol* 77:586–596. <https://doi.org/10.1128/AEM.01080-10>.
- Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, Dewinter A, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Vieceli J, Wegener J, Wu D, Yang A, Zaccarin D, Zhao P, Zhong F, Korlach J, Turner S. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* 323:133–138. <https://doi.org/10.1126/science.1162986>.
- Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
- Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpidis NC. 2010. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 7:455–457. <https://doi.org/10.1038/nmeth.1457>.
- Pruesse E, Quast C, Knittel K, Fuchs BM, Ludwig W, Peplies J, Glöckner FO. 2007. SILVA: a comprehensive online resource for quality checked and aligned ribosomal RNA sequence data compatible with ARB. *Nucleic Acids Res* 35.21:7188–7196.

12. Markowitz VM, Mavromatis K, Ivanova NN, Chen I-MA, Chu K, Kyrpides NC. 2009. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 25:2271–2278. <https://doi.org/10.1093/bioinformatics/btp393>.
13. Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, Dehal P, Ware D, Perez F, Canon S, Sneddon MW, Henderson ML, Riehl WJ, Murphy-Olson D, Chan SY, Kamimura RT, Kumari S, Drake MM, Brettin TS, Glass EM, Chivian D, Gunter D, Weston DJ, Allen BH, Baumohl J, Best AA, Bowen B, Brenner SE, Bun CC, Candonia J-M, Chia J-M, Colasanti R, Conrad N, Davis JJ, Davison BH, DeJongh M, Devoid S, Deitrich E, Dubchak I, Edirisnghe JN, Fang G, Faria JP, Frybarger PM, Gerlach W, Gerstein M, Greiner A, Gurtowski J, Haun HL, He F, Jain R, Joachimiak MP, Keegan KP, Kondo S, Kumar V, Land ML, Meyer F, Mills M, Novichkov PS, Oh T, Olsen GJ, Olson R, Parrello B, Pasternak S, Pearson E, Poon SP, Price GA, Ramakrishnan S, Ranjan P, Ronald PC, Schatz MC, Seaver SMD, Shukla M, Sutormin RA, Syed MH, Thomason J, Tintle NL, Wang D, Xia F, Yoo H, Yoo S, Yu D. 2018. KBase: the United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* 36:566–569. <https://doi.org/10.1038/nbt.4163>.
14. Ondov BD, Treangen TJ, Melsted P, Mallonee AB, Bergman NH, Koren S, Phillippy AM. 2016. Mash: fast genome and metagenome distance estimation using MinHash. *Genome Biol* 17:132. <https://doi.org/10.1186/s13059-016-0997-x>.
15. Yin Y, Mao X, Yang J, Chen X, Mao F, Xu Y. 2012. dbCAN: a Web resource for automated carbohydrate-active enzyme annotation. *Nucleic Acids Res* 40:W445–W451. <https://doi.org/10.1093/nar/gks479>.
16. Hwang DM, Dempsey A, Tan K-T, Liew C-C. 1996. A modular domain of NifU, a nitrogen fixation cluster protein, is highly conserved in evolution. *J Mol Evol* 43:536–540. <https://doi.org/10.1007/PL00006114>.
17. Gtari M, Ghodhbane-Gtari F, Nouioui I, Beauchemin N, Tisa LS. 2012. Phylogenetic perspectives of nitrogen-fixing actinobacteria. *Arch Microbiol* 194:3–11. <https://doi.org/10.1007/s00203-011-0733-6>.